

References

10777 Gd-IgA1 (KM55) Anti-Human Rat IgG MoAb - IBL

No.	Field	Keywords	Published	Title
1	Dermatology	Nephritis, Vasculitis	2019	Galactose deficient IgA1 (GD-IgA1) in skin and serum from patients with skin-limited and systemic IgA Vasculitis. Neufeld M et al. J Am Acad Dermatol. 2019 Mar 19; pii: S0190-9622(19)30443-8.
2	Nephrology	MEST-C, Herramientas	2021	Measurement of galactosyl-deficient IgA1 by the monoclonal antibody KM55 contributes to predicting patients with IgA nephropathy with high risk of long-term progression. Martín-Penagos L et al. Nefrologia. May-Jun 2021;41(3):311-320.
3	Nephrology	Alport syndrom	2021	Utility of glomerular Gd-IgA1 staining for indistinguishable cases of IgA nephropathy or Alport syndrome. Ishiko S et al. Clin Exp Nephrol. 2021 Mar 20.
4	Nephrology	KM55 monoclonal antibody	2021	KM55 Monoclonal Antibody Staining in IgA-Dominant Infection-Related Glomerulonephritis. Zhang M et al. Nephron. 2021;145(3):225-237.
5	Nephrology	Renal involvement	2021	Glomerulonephritis with severe nephrotic syndrome induced by immune complexes composed of galactose-deficient IgA1 in primary Sjögren's syndrome: a case report. Nishioka R et al. BMC Nephrol. 2021 Mar 25;22(1):108.
6	Nephrology	Biomarker, Immune complexes	2019	Biomarkers for IgA nephropathy on the basis of multi-hit pathogenesis. Suzuki H. Clin Exp Nephrol. 2019 Jan;23(1):26-31.
7	Nephrology	Epitope	2018	Galactose-Deficient IgA1-Specific Antibody Recognizes GalNAc-Modified Unique Epitope on Hinge Region of IgA1. Yamasaki K et al. Monclon Antib Immunodiagn Immunother. 2018 Dec;37(6):252-256.
8	Nephrology	HAA	2015	Novel lectin-independent approach to detect galactose-deficient IgA1 in IgA nephropathy. Yasutake J et al. Nephrol Dial Transplant. 2015 Aug;30(8):1315-21.
9	Nephropathy	Complement activation, KM55 antibody	2019	Clinical Significance of Galactose-Deficient IgA1 by KM55 in Patients with IgA Nephropathy. Zhang K et al. Kidney Blood Press Res. 2019;44(5):1196-1206.
10	Nephropathy	Mesangial- Gd-IgA1	2018	Clinical significance of serum and mesangial galactose-deficient IgA1 in patients with IgA nephropathy. Wada Y et al. PLoS One. 2018 Nov 2;13(11):e0206865.
11	Nephropathy	Vasculitis, Renal biopsy	2012	IgA nephropathy and IgA vasculitis with nephritis have a shared feature involving galactose-deficient IgA1-oriented pathogenesis. Suzuki H et al. Kidney Int. 2018 Mar;93(3):700-705.

Distributed by:



Immuno-Biological Laboratories, Inc.
8201 Central Ave NE, Suite P
Minneapolis, MN 55432

Toll-Free: 888-523-1246
Email: info@IBL-America.com
Web: www.IBL-America.com