



Immuno-Biological Laboratories Co., Ltd.

References

[27201 GIP Active Form \(Human\) Assay Kit - IBL](#)

No.	Field	Keywords	Published	Title
1	Metabolic	Canagliflozin, Type 2 diabetes	2017	Effect of canagliflozin on circulating active GLP-1 levels in patients with type 2 diabetes: a randomized trial. Takebayashi K et al. Endocr J. 2017 Sep 30;64(9):923-931.
2	Metabolic	Diabetes, Glucagon	2020	Plasma Glucagon Levels Measured by Sandwich ELISA Are Correlated With Impaired Glucose Tolerance in Type 2 Diabetes. Kobayashi M et al. Endocr J. 2020;10.1507/endocrj.EJ20-0079.
3	Metabolic	Dipeptidyl, Receptor-mediated	2016	Dipeptidyl peptidase-4 inhibitor treatment induces a greater increase in plasma levels of bioactive GIP than GLP-1 in non-diabetic subjects. Yanagimachi T et al. Mol Metab. 2016 Dec 31;6(2):226-231.
4	Metabolic	MAGE, Type 2 diabetes	2014	Teneligliptin improves glycemic control with the reduction of postprandial insulin requirement in Japanese diabetic patients. Tsuchimochi W et al. Endocr J. 2014 Sep 25.
5	Metabolic	Type 2 diabetes	2014	Addition of sitagliptin or metformin to insulin monotherapy improves blood glucose control via different effects on insulin..... Otsuka Y et al. Endocr J. 2014 Oct 19.
6	Neurology	Neuro- inflammation	2016	Glucose-Dependent Insulinotropic Polypeptide Ameliorates Mild Traumatic Brain... Yu YW et al. J Neurotrauma. 2016 Nov 15;33(22):2044-2054. Epub 2016 May 9.
7	Other	Pancreatectomy	2020	Pseudo-hyperglucagonemia Was Observed in the Pancreatectomized Cases When Measured by Glucagon Sandwich ELISA. Kobayashi M et al. J Diabetes Investig. 2020;10.1111/jdi.13325.

Distributed by:



Immuno-Biological Laboratories, Inc.
8201 Central Ave NE, Suite P
Minneapolis, MN 55432

Toll-Free: 888-523-1246
Email: info@IBL-America.com
Web: www.IBL-America.com