

References

27412 Angiotensinogen (Total) Assay Kit - IBL

No.	Field	Keywords	Published	Title
1	Cardiovascular	Blood pressure, African Ancestry	2014	Urinary angiotensinogen excretion is associated with blood pressure independent of the circulating renin-angiotensin system in a group of african ancestry. Michel FS et. al. Hypertension. 2014 Jul;64(1):149-56.
2	Cardiovascular	Blood pressure, UAGT	2010	Urinary angiotensinogen is correlated with blood pressure in men (Bogalusa Heart Study) Kobori H et al. J Hypertens. 2010 Jul;28(7):1422-8.
3	Cardiovascular	Diabetes, CKD	2019	Urinary angiotensinogen antedates the development of stage 3 CKD in patients with type 1 diabetes mellitus. Ba Aqeel S et al. Physiol Rep. 2019 Oct;7(19):e14242.
4	Cardiovascular	Left ventricular mass index	2015	Association between urinary angiotensinogen excretion rates and left ventricular mass index and carotid intima-media thickness....Tiryaki O et al. Clin Transplant. 2015 Apr;29(4):351-8.
5	Renal Disease	AKI, CKD	2019	The Role of Urinary Angiotensinogen in Kidney Interstitial Inflammation and Renal Prognosis. Dou Y et al. Iran J Kidney Dis. 2019 Nov;13(6):372-379.
6	Renal Disease	Blood pressure, Plasma	2008	Urinary angiotensinogen as a potential biomarker of severity of chronic kidney diseases. Kobori H et al. J Am Soc Hypertens. 2008 Sep-Oct;2(5):349-54.
7	Renal Disease	CKD, Circadian rhythm	2014	Circadian rhythm of plasma and urinary angiotensinogen in healthy volunteers and in patients.... Nishijima Y et al. J Renin Angiotensin Aldosterone Syst. 2014 Dec;15(4):505-8.
8	Renal Disease	Glomerulonephritis IgAN	2014	Changes in urinary angiotensinogen posttreatment in pediatric IgA nephropathy patients. Urushihara M et al. Pediatr Nephrol. 2014 Dec 20.
9	Renal Disease	Urinary tract, Cicatrix	2015	Urinary Biomarkers for Screening for Renal Scarring in Children with Febrile Urinary Tract Infection: Pilot Study. Kitao T et al. J Urol. 2015 Apr 29. pii: S0022-5347(15)03891-4.
10	Other	ANG, hAGT	2007	Novel sandwich ELISA for human angiotensinogen. Katsurada A et al. Am J Physiol Renal Physiol. 2007 Sep;293(3):F956-60.
11	Other	Preterm neonate RAS	2015	Urinary angiotensinogen level is increased in preterm neonates. Suzue M et al. Clin Exp Nephrol. 2015 Apr;19(2):293-7.
12	Other	Type 2 diabetes	2014	Poor glycemic control and decreased renal function are associated with increased intrarenal RAS activity.... Nakatani S et al. Diabetes Res Clin Pract. 2014 Jul;105(1):40-6.

Distributed by:



Immuno-Biological Laboratories, Inc.
8201 Central Ave NE, Suite P
Minneapolis, MN 55432

Toll-Free: 888-523-1246
Email: info@IBL-America.com
Web: www.IBL-America.com