



References

[27782 Soluble \(Pro\)renin Receptor Assay Kit - IBL](#)

No.	Field	Keywords	Published	Title
1	Blood Pressure	d ose P	2019	Losartan prevents the elevation of blood pressure in adipose-PRR deficient female mice while elevated circulating sPRR activates the renin-angiotensin system. Gatineau E et al. Am J Physiol Heart Circ Physiol. 2019 Mar 1;316(3):H506-H515.
2	Inflammatory/Ocular	RAPS, (P)RR	2017	A Novel Single-Strand RNAi Therapeutic Agent Targeting the (Pro)renin Receptor Suppresses Ocular... Kanda A et al. Mol Ther Nucleic Acids. 2017 Jun 16;7:116-126.
3	Metabolic	Diabetes,	2019	Urinary angiotensinogen antedates the development of stage 3 CKD in patients with type 1 diabetes mellitus. Ba Aqeel S et al. Physiol Rep. 2019 Oct;7(19):e14242.
4	Pancreatic	Left ventricular	2015	Association between urinary angiotensinogen excretion rates and left ventricular mass index and carotid intima-media thickness....Tiryaki O et al. Clin Transplant. 2015 Apr;29(4):351-8.
5	Plasma	AKI,	2019	The Role of Urinary Angiotensinogen in Kidney Interstitial Inflammation and Renal Prognosis. Dou Y et al. Iran J Kidney Dis. 2019 Nov;13(6):372-379.
6	Plasma	Blood pressure,	2008	Urinary angiotensinogen as a potential biomarker of severity of chronic kidney diseases. Kobori H et al. J Am Soc Hypertens. 2008 Sep-Oct;2(5):349-54.
7	Prenatal	CKD,	2014	Circadian rhythm of plasma and urinary angiotensinogen in healthy volunteers and in patients.... Nishijima Y et al. J Renin Angiotensin Aldosterone Syst. 2014 Dec;15(4):505-8.
8	Prenatal	Glomerulonephritis IgAN	2014	Changes in urinary angiotensinogen posttreatment in pediatric IgA nephropathy patients. Urushihara M et al. Pediatr Nephrol. 2014 Dec 20.
9	Prenatal	Urinary tract,	2015	Urinary Biomarkers for Screening for Renal Scarring in Children with Febrile Urinary Tract Infection: Pilot Study. Kitao T et al. J Urol. 2015 Apr 29. pii: S0022-5347(15)03891-4.
10	Renal Disease	ANG,	2007	Novel sandwich ELISA for human angiotensinogen. Katsurada A et al. Am J Physiol Renal Physiol. 2007 Sep;293(3):F956-60.
11	Renal Disease	Preterm neonate	2015	Urinary angiotensinogen level is increased in preterm neonates. Suzue M et al. Clin Exp Nephrol. 2015 Apr;19(2):293-7.
12	Serum	Type 2 diabetes	2014	Poor glycemic control and decreased renal function are associated with increased intrarenal RAS activity.... Nakatani S et al. Diabetes Res Clin Pract. 2014 Jul;105(1):40-6.

Distributed by:



Immuno-Biological Laboratories, Inc.
8201 Central Ave NE, Suite P
Minneapolis, MN 55432

Toll-Free: 888-523-1246
Email: info@IBL-America.com
Web: www.IBL-America.com