



Mac2-bp

NASH (Non-Alcoholic Steatohepatitis)

ELISA (96Well)


- Research Use Only-

Mac-2 binding protein (Mac-2bp), known as 90K, is a highly N-glycosylated, secreted protein, identified as a ligand of Galectin-3. It is considered that through interaction with Galectin-3, Mac-2bp promotes homotypic cell-cell contact or regulates cell adhesion. And it has been reported that Mac-2bp levels in blood have associations with various cancers and chronic hepatic diseases such as NASH (Non-Alcoholic Steatohepatitis).

Product Code	Sample Type	Product Name	Measuring Range	Measuring Samples
27362	Human	Human Mac-2 binding protein (Mac-2bp) Assay Kit - IBL	0.78 - 100 ng/mL	Serum, EDTA-Plasma, Cell culture supernatant
27796	Mouse	Mouse Mac-2 binding protein (Mac-2bp) Assay Kit - IBL	0.78 - 50 ng/mL	Serum, EDTA-Plasma, Tissue extract, Cell culture supernatant

IBL

Human Mac-2 binding protein Assay Kit - IBL



Human Mac-2 binding protein Assay Kit - IBL Code No. 27362

27362 Human Mac-2 binding protein (Mac-2bp) Assay Kit - IBL

Reference

A novel noninvasive diagnostic method for nonalcoholic steatohepatitis using two glycobiomarkers. Kamada Y et al. Hepatology. 2015 Jul 21.


Serum Mac-2 binding protein levels as a novel diagnostic biomarker for prediction of disease severity and nonalcoholic steatohepatitis. Kamada Y et al. Proteomics Clin Appl. 2013 Jun 14.

Serum Mac-2 binding protein is a novel biomarker for chronic pancreatitis. Maekawa T et al. World J Gastroenterol. 2016 May 7;22(17):4403-10.

27796 Mouse Mac-2 binding protein (Mac-2bp) Assay Kit - IBL

Reference

Establishment of mouse Mac-2 binding protein enzyme-linked immunosorbent assay and its application for mouse chronic liver disease models. Iwata A et al. Hepatol Res. 2016 Sep 11.



Explanation of Method of Feeding

Normal Meal: The meal that is used as a normal meal at an animal testing center was fed to mice.

NASH Meal: High fat and high cholesterol food (7.5% fatty acid, 1.25% cholesterol, 0.5% and 0.5% cholic acid) was fed to mice for 4 weeks or 8 weeks.

Data provided by: Dr. Kamata Y and Dr. Miyoshi Y, Department of Functional Diagnostic Science, Osaka University, Graduate School of Medicine

Distributed by:



Immuno-Biological Laboratories, Inc.
8201 Central Ave NE, Suite P
Minneapolis, MN 55432

Toll-Free: 888-523-1246
Email: info@IBL-America.com
Web: www.IBL-America.com