



## PRODUCT INFORMATION

### Nickel HRP Conjugate – 1.0 mg

PRODUCT CODE: X-CON-0010-1MG

STORAGE: 2 - 8 °C, protected from sun light.

### PRODUCT DESCRIPTION

Metal affinity interaction allows polyhistidine tag containing protein to be complexed and detected with molecules that contain chelated divalent ions such as  $\text{Ni}^{2+}$ ,  $\text{Cu}^{2+}$ ,  $\text{Co}^{2+}$  and  $\text{Zn}^{2+}$ . Horseradish peroxidase (HRP) oxidizes corresponding substrates with high efficiency, generating colorimetric or chemiluminescent reactions and is frequently used as a reporter enzyme for sensitive assays like ELISA, immunohistochemistry and western blot. HRP is conjugated with chelated  $\text{Ni}^{2+}$  under optimal conditions. Nickel HRP Conjugate is useful as a reagent for detecting polyhistidine tag containing proteins in ELISA and western-blotting procedures.

### PRECAUTIONS AND DISCLAIMER

This product is for R&D use only, not for drug, household, or other uses. Please consult the Material Safety Data Sheet for information regarding hazards and safe handling practices.

### FORMULATION

For shipping at ambient temperature Nickel HRP Conjugate is dried with a HEPES, NaCl, sucrose buffer base.

### PREPARATION AND HANDLING

The product should be reconstituted with 100  $\mu\text{l}$  water yielding a concentration of 10 mg/ml. The reconstituted stock solution can be frozen in aliquots for later usage. Stock solutions can be diluted in buffers containing > 0.1 % BSA as needed. Avoid using buffer that contain EDTA or other metal chelators, avoid reducing agents, sodium acid and imidazole.

### STORAGE / STABILITY

For long term storage the dry-stabilized Nickel HRP Conjugate should be stored between 2 °C and 8 °C. Reconstituted stock solutions can be stored at 2 - 8 °C for up to 2 weeks. For long term storage, stock solutions can be frozen in working aliquots. Repeated freeze-thaw cycles should be avoided.

### RECONSTITUTION AND CONCENTRATION

10 mg/ml after reconstitution with 100  $\mu\text{l}$   $\text{H}_2\text{O}$ .

### RECOMMENDED ELISA DILUTION

1:500 – 1: 5000 in secondary ELISA detection. For optimal performance the reagent should be titrated for each application.

### RECOMMENDED RETEST DATE

09/2022

### BACKGROUND REFERENCES

1. Wong, J., et al., Direct force measurements of the streptavidin –biotin interaction, *Biomolecular Engineering*, 16, 45-55 (1999).
2. Hochuli, E., et al., New metal chelate adsorbent for proteins and peptides containing neighbouring histidine residues, *J. Chromatogr.*, 411, 177-84 (1987).



Distributed By:  
**IBL-America, Inc.**  
 8201 Central Ave NE, Suite P  
 Minneapolis, MN 55432, USA  
[info@ibl-america.com](mailto:info@ibl-america.com)  
 (888) 523 1246