

Code No. 10097

Anti-S-100 Protein (2A10) Mouse Ascites MoAb**Volume** : 0.5ml**Lot:**

Introduction : S100 proteins (16 members) show a very divergent pattern of cell- and tissue-specific expression, of subcellular localizations and relocations, of post-translational modifications, and of affinities for Ca²⁺, Zn²⁺, and Cu²⁺, consistent with their pleiotropic intra- and extracellular functions. Up to 40 target proteins are reported to interact with S100 proteins and for S100A1 alone 15 target proteins are presently known. Therefore it is not surprising that many functional roles have been proposed and that several human disorders such as cancer, neurodegenerative diseases, cardiomyopathies, inflammations, diabetes, and allergies are associated with an altered expression of S100 proteins.

Antigen : Bovine Brain S-100 Protein**Source** : Mouse-Mouse hybridoma**Clone** : 2A10**Subclass** : IgG1**Purification** : Affinity purified with Protein A**Form** : Lyophilized product from 1% BSA in PBS containing 0.05%NaN₃**How to use** : 0.5 ml distilled water will be added to the product**Dilution** : PBS (pH7.4) containing 1% BSA**Stability** : Lyophilized product, 5 years at 2 – 8 °C
: Solution, 2 years at –20 °C

Application : This antibody can be stained in formalin fixed paraffin embedded tissues by several Immunohistochemical techniques such as Avidin Bition Complex (ABC) Method. The optimal dilution is X 200, however, the dilution rate should be optimized by each laboratories.

Specificity : Reacts with β chain of Bovine and Human

For research use only, not for use in diagnostic procedures.

Distributed by:



Immuno-Biological Laboratories, Inc. Toll-Free: 888-523-1246

8201 Central Ave NE, Suite P

Minneapolis, MN 55432

Email: info@IBL-America.comWeb: www.IBL-America.com