

Code No. 10907

**Anti-
CD25 (H-48) Mouse IgG MoAb**

Volume : 100 µg

-
- Introduction** : CD25 is IL-2 receptor α chain (IL-2R α which is a low-affinity receptor of IL-2. CD25 forms high-affinity IL-2 receptors associating with CD122 (β chain) and CD132 (γ chain). CD25 is expressed in activated T cells and B cells. Increase of CD25 in serum is observed in individuals suffer adult T cell leukemia or malignant lymphoma.
- Antigen** : Purified human IL-2 receptor α
- Source** : Mouse-Mouse hybridoma
(SP2/0 \times BALB/c mouse spleen cells)
- Clone** : H-48
- Subclass** : IgG1
- Purification** : Gel filtration chromatography following ammonium sulfate precipitation
- Form** : Lyophilized product from PBS containing 1 % BSA and 0.05 % NaN₃
- How to use** : 0.2 mL deionized water will be added to the product, then its concentration comes to 500 µg/mL
- Stability** : Lyophilized product, 5 years at 2 - 8 °C
: Solution, 2 years at -20 °C
- Application** : This antibody can be used for western blotting in concentration of 10 µg/mL.
: This antibody can be used for immuno-precipitation at 10 µg /test
: This antibody can be used for flow cytometry analysis at 1 µg (per 1 \times 10⁵ cells)
: This antibody can be used for ELISA (as a capture antibody) in concentration of 5 µg/mL
- Reference** : Tanaka Y, Inoi T, Tozawa H, Sugamura K, Hinuma Y. New monoclonal antibodies that define multiple epitopes and a human-specific marker on the interleukin 2 receptor molecules of primates. *Microbiol Immunol.* 1986;30(4):373-88.

For research use only, not for use in diagnostic procedures.

Distributed by:

Immuno-Biological Laboratories, Inc.
8201 Central Ave NE, Suite P
Minneapolis, MN 55432Toll-Free: 888-523-1246
Email: info@IBL-America.com
Web: www.IBL-America.com

Code No. 10907

**Anti-
CD25 (H-48) Mouse IgG MoAb**

Volume : 500 µg

Introduction : CD25 is IL-2 receptor α chain (IL-2R α which is a low-affinity receptor of IL-2. CD25 forms high-affinity IL-2 receptors associating with CD122 (β chain) and CD132 (γ chain). CD25 is expressed in activated T cells and B cells. Increase of CD25 in serum is observed in individuals suffer adult T cell leukemia or malignant lymphoma.**Antigen** : Purified human IL-2 receptor α **Source** : Mouse-Mouse hybridoma
(SP2/0 \times BALB/c mouse spleen cells)**Clone** : H-48**Subclass** : IgG1**Purification** : Gel filtration chromatography following ammonium sulfate precipitation**Form** : Lyophilized product from PBS containing 1 % BSA and 0.05 % NaN₃**How to use** : 1.0 mL deionized water will be added to the product, then its concentration comes to 500 µg/mL**Stability** : Lyophilized product, 5 years at 2 - 8 °C
: Solution, 2 years at -20 °C**Application** : This antibody can be used for western blotting in concentration of 10 µg/mL.
: This antibody can be used for immuno-precipitation at 10 µg /test
: This antibody can be used for flow cytometry analysis at 1 µg (per 1 \times 10⁵ cells)
: This antibody can be used for ELISA (as a capture antibody) in concentration of 5 µg/mL**Reference** : Tanaka Y, Inoi T, Tozawa H, Sugamura K, Hinuma Y. New monoclonal antibodies that define multiple epitopes and a human-specific marker on the interleukin 2 receptor molecules of primates. Microbiol Immunol. 1986;30(4):373-88.

For research use only, not for use in diagnostic procedures.

Distributed by:

Immuno-Biological Laboratories, Inc.
8201 Central Ave NE, Suite P
Minneapolis, MN 55432Toll-Free: 888-523-1246
Email: info@IBL-America.com
Web: www.IBL-America.com