

Code No. 10915

**Anti-
CD252/OX40L (TAG34) Mouse IgG MoAb**

Volume : 100 µg

Introduction : CD252/OX40L is a molecule identical to gp34 expressed in T cells infected with HTLV-1 and it is expressed on activated B cell, dendritic cells and vascular endothelial cells as a trimer belonging to TNF superfamily. CD252/OX40L have a role in proliferation of T cells and production of cytokines through OX40 expressed on activated T cells. In addition, OX40L expressed on the activated vascular endothelial cells are involved in activated T cell invasion to inflamed area.

Antigen : Lysate of HTLV-1 persistently-infected cell line (MT-2)

Source : Mouse-Mouse hybridoma
(SP2/0 × BALB/c mouse spleen cells)

Clone : TAG34 **Subclass** : IgG₁

Purification : Gel filtration chromatography following ammonium sulfate precipitation

Form : Lyophilized product from PBS containing 1 % BSA and 0.05 % NaN₃

How to use : 1.0 mL deionized water will be added to the product, then its concentration comes to 100 µg/mL

Stability : Lyophilized product, 5 years at 2 - 8 °C
: Solution, 2 years at -20 °C

Application : This antibody can be used for western blotting in concentration of 10 µg/mL.
: This antibody can be used for immuno-precipitation at 10 µg /test.
: This antibody can be used for flow cytometry analysis at 1 µg (per 1×10⁵ cells)
: This antibody can be used for blocking in concentration of 20 µg/mL

Reference : 1. Takahashi Y, Tanaka Y, Yamashita A, Koyanagi Y, Nakamura M, Yamamoto N. OX40 stimulation by gp34/OX40 ligand enhances productive human immunodeficiency virus type 1 infection. *J Virol.* 2001 Aug;75(15):6748-57.
2. Tozawa H, Andoh S, Takayama Y, Tanaka Y, Lee B, Nakamura H, Hayami M, Hinuma Y. Species-dependent antigenicity of the 34-kDa glycoprotein found on the membrane of various primate lymphocytes transformed by human T-cell leukemia virus type-I (HTLV-I) and simian T-cell leukemia virus (STLV-I). *Int J Cancer.* 1988 Feb 15;41(2):231-8.
3. Tanaka Y, Inoi T, Tozawa H, Yamamoto N, Hinuma Y. A glycoprotein antigen detected with new monoclonal antibodies on the surface of human lymphocytes infected with human T-cell leukemia virus type-I (HTLV-I). *Int J Cancer.* 1985 Nov 15;36(5):549-55.

For research use only, not for use in diagnostic procedures.

Distributed by:



Immuno-Biological Laboratories, Inc. Toll-Free: 888-523-1246

8201 Central Ave NE, Suite P

Minneapolis, MN 55432

Email: info@IBL-America.comWeb: www.IBL-America.com