

Code No. 18111

Anti-Human SCF (K089) Rabbit IgG Affinity Purify

Volume : 100 µg

Introduction: Stem cell factor (SCF) and its receptor KIT play an important role in various

biologic phases, such as hematopoiesis, reproduction, and regeneration. Structurally, c-Kit contains five immunoglobulin-like domains extracellularly and a catalytic domain divided into two regions by a 77 amino acid insert intracellularly. Studies in white spotting and steel mice have shown that functional SCF and c-Kit are critical in the survival and development of stem cells involved in hematopoiesis, pigmentation and reproduction. Mutations in c-Kit are associated with a variety of human diseases. Interaction of SCF with c-Kit rapidly induces receptor dimerization and increases in autophosphorylation activity. Downstream of c-Kit, transduction components multiple are activated. including phosphatidylinositol-3-kinase, Src family members, the JAK/STAT pathway and

the Ras-Raf-MAP kinase cascade.

: Synthetic peptide of a part of Human SCF Antigen

Purification: Purified with antigen peptide

: Lyophilized product from 1 % BSA in PBS containing 0.05 % NaN₃ **Form**

How to use : 1.0 mL deionized water will be added to the product (the conc. comes up 100 μg /mL)

Stability : Lyophilized product, 5 years at 2 – 8 °C

: Solution, 2 years at -20 °C

Application: This antibody can be used for immunohistochemistry with formalin fixed paraffin

embedded tissues by several techniques such as Avidin Biotin Complex (ABC) Method. The optimal concentration is 1-5 µg/mL, however, the concentration should

be optimized by each laboratory.

: This antibody can be used for western blotting in concentration of 1-5 µg /mL.

Specificity: Human SCF specific.