

Code No. 18281

Anti-Rat

HGFα (H56) Rabbit IgG Affinity Purify

Volume : 100 µg

Introduction: Hepatocyte Growth factor (HGF) was discovered as a mitogen for hepatocytes.

HGF was subsequently found to be identical to the scatter factor, which destroys epithelial cell adhesion and promotes cell movement. Some reports have Hepatocyte Growth factor (HGF) was discovered as a mitogen for hepatocytes. HGF was subsequently found to be identical to the scatter factor, which destroys epithelial cell adhesion and promotes cell movement. Some reports have shown that HGF is expressed in normal and malignant mammary epithelium. HGF has also been reported to promote motility and growth of epithelial cells, to induce morphogenesis of epithelial cells and to promote vascularization. It has been speculated that HGF is involved in the growth and metastasis of cancer cells. The first step in the initiation of HGF action is dependent on its biding to a specific cell surface receptor, the HGF receptor, encoded by the proto-oncogene c-met. It has been suggested that c-Met mediates both responses, i.e., promotion of growth and motility of HGF.

HGF is synthesized as a 728 amino acid that is processed to generate the mature

growth factor consisting of a disulfide-linked 69 kDa α 34 kDa β chain.

: Synthetic peptide for N-Terminal of rat HGF **Antigen**

(PLVKIKTKKVNSADEC)

Purification: Purified with antigen peptide

Form : Lyophilized product from PBS containing 1% BSA and 0.05% NaN₃

How to use : 1.0 mL deionized water will be added to the product (the conc. comes up 100 μg/mL)

Stability : Lyophilized product, 5 years at 2 - 8 °C

: Solution, 2 years at -20 °C

Application: This antibody can be used in immunohistchemistry with formalin fixed paraffin

embedded tissues after trypsin treatment, by several techniques such as Avidin Biotin Complex (ABC) Method. The optimal concentration is 2-5 µg/mL, however,

concentration should be optimized by each laboratory.

: This antibody can be used for western blotting in concentration of 2-5 µg /mL.

: Rat HGF-α specific. Specificity

Not cross-react with Human HGF and Rat HGFβ.

For research use only, not for use in diagnostic procedures.



Immuno-Biological Laboratories, Inc. Toll-Free: 888-523-1246 8201 Central Ave NE, Suite P Minneapolis, MN 55432

Email: info@IBL-America.com Web: www.IBL-America.com