

Code No. 18420

Anti-Rat

VEGF (V-N) Rabbit IgG Affinity Purify

Volume : 100 µg

Introduction: Vascular Endothelial Cell Growth Factor (VEGF) is a homodimeric protein initially purified from media conditioned by normal bovine pituitary folliculo-stellate cells and secreted by a variety of vascularized tissues. It was subsequently found to be identical to a vascular permeability factor (VPF), which was previously identified in media conditioned by tumor cell lines based upon its ability to increase the permeability of capillary blood vessels. The reported activities of VEGF include stimulation of endothelial cell growth, angiogenesis and capillary permeability. Human VEGF is a 38.2 kDa homodimeric protein consisting of two 165 amino acid polypeptide chains. VEGF is expressed in many human tumor cells, including human adenocarcinoma, human pancreatic carcinoma, human hepatocellular carcinoma, renal cell carcinoma, fibrosarcoma, HL60 promyelocytic leukemia, GS-9L glioma and U937 lymphoma cells. In normal tissues, VEGF expression has been observed in activated macrophages, keratinocytes, hepatocytes, smooth muscle cells Leydig cells, embryonic fibroblasts and bronchial and choroids plexus epithelium, renal glomerular visceral epithelium and mesangial cells.

Antigen : Synthetic peptide of the N terminal part of Rat VEGF

Purification: Purified with antigen peptide

: Lyophilized product from 1 % BSA in PBS containing 0.05% NaN₃ **Form**

How to use : 1.0 mL deionized water will be added to the product (the conc. comes up 100 μg/mL)

Stability : Lyophilized product, 5 years at 2 – 8 °C

: Solution, 2 years at -20 °C

Application: This antibody can be used for western blotting in concentration of 1 - 5 µg /mL.

Specificity: Reacts with both of reduced form and non-reduced form.

For research use only, not for use in diagnostic procedures.



