

Code No. 18669

Anti-Human

IκB-α (1037) Rabbit IgG Affinity Purify

Volume : $100 \, \mu g$

Lot No

Introduction : In signal transduction system, IkB acts as inhibitor to forms complex with

> NF-κB in cytoplasm and subsequently to inactivate NF-κB. NF-κB is transcription factor distributed in various cells. NF-kB is considered to control various gene expression by degradation of IkB and subsequent translocation of degradation products to nuclear in accordance with various

stimulation.

Antigen : Synthetic peptide in portion of N-terminus of Human $I\kappa B-\alpha$ protein

Purification : Purified with antigen peptide

Form : Lyophilized product from 1% BSA in PBS containing 0.05% NaN₃

: 1.0 ml distilled water will be added to the product How to use

Dilution : PBS (pH7.4) containing 1% BSA

Stability : Lyophilized product, 5 years at 2 – 8 °C

: Solution, 2 years at -20 °C

Application : This antibody can be used for western blotting in concentration of

 $1~2 \mu g/ml$.

Specificity : Confirmed by Human IκB-α

> The sequence of synthetic peptide used for antigen has homology of 100% to that of mouse. However, sample of mouse by this antibody is not tested.

> > For research use only, not for use in diagnostic procedures.

