

Code No. 18671

Anti-Human

*α*Synuclein (S122) Rabbit IgG Affinity Purify

Volume : $100 \, \mu g$

: The pre-synaptic protein, α -synuclein, has been associated with the

pathogenesis of several neurodegenerative diseases. These include prevalent conditions such as Parkinson's disease (PD), dementia with Lewy bodies (DLB), and the Lewy body variant of Alzheimer's disease (LBVAD), as well as

rarer conditions including multiple systems atrophy (MSA), and

neurodegeneration with brain iron accumulation type-1 (NBIA-1). Tha

antibody may be useful for these research.

Antigen : Synthetic peptide for C-terminal of Humanα-Synuclein

(DNEAYEMPSEEGYQDY)

Purification : Affinity Purified with synthetic peptide

Form : Lyophilized product from 1% BSA in PBS containing 0.05%NaN₃

How to use : 1.0 ml distilled water will be added to the product

Dilution : PBS (pH7.4) containing 1% BSA

Stability : Lyophilized product, 5 years at 2 – 8 °C

: Solution, 2 years at -20 °C

Application : This antibody can be stained in formalin fixed paraffin embedded tissues after

> microwave treatment by several Immunohistochemical techniques such as Avidin Bition Complex (ABC) Method. The optimal dilution is about 2 µg/ml,

however, the dilution rate should be optimized by each laboratories.

: This antibody can be used for western blotting in concentration of $0.5 \sim 2 \mu g$ /ml.

Specificity : Confirmed by western blotting

Reactivity : Cross reacts with mouse and rat

References : Ueda K. et. al. Molecular cloning of cDNA encoding an unrecognized

component of amyloid in Alzheimer disease. Proc. Natl. Acad. Sci. U.S.A.

1993: **90** (23), 11282- 11286

Masliah E. et. al. Dopaminergic loss and inclusion body formation in α

Synuclein mice: implications for neuro- degenerative disorders. Science 2000:

287, 1265-1269

For research use only, not for use in diagnostic procedures.

Distributed by:



Email: info@IBL-America.com Web: www.IBL-America.com