

Code No. 18957

**Anti-Human
sAPP β -Wild Type Rabbit IgG Affinity Purify**Volume : 100 μ g
Lot No : 0B-521

-
- Introduction** : Amyloid precursor protein (APP) is precursor protein of Amyloid β which is major constituent of senile plaque in Alzheimer's disease. It is known that there are three major isoforms, APP695, APP751 and APP770, and are generated from alternative splicing of common precursor mRNA. Processing of APP occurs by two major pathways, non-amyloidogenic pathway and amyloidogenic pathway. The non-amyloidogenic pathway is mediated by α and γ -secretases and gives rise to a large fragment known as soluble APP α (sAPP α) and a small 3 kDa peptide known as p3. On the other hand, the Amyloidogenic pathway is mediated by β - and γ -secretases and yields soluble APP β (sAPP β) and Amyloid β . The physiologic function of APP itself is not clear, however, it is supposed that the function of APP in neuron system is different from that in other organ.
- Antigen** : Synthetic peptide in portion of C-terminus of Human sAPP β -Wild Type (ISEVKM)
- Purification** : Purified with antigen peptide
- Form** : Lyophilized product from 1% BSA in PBS containing 0.05% NaN₃
- How to use** : 1.0 ml distilled water will be added to the product, then its concentration comes to 100 μ g/ml
- Dilution** : PBS (pH7.4) containing 1% BSA
- Stability** : Lyophilized product, 5 years at 2 – 8
: Solution, 2 years at –20
- Application** : This antibody can be used for western blotting in concentration of about 2 μ g/ml.
: This antibody can be used for immunoprecipitation in concentration of about 3 μ g/ml.
- Specificity** : This antibody can detect sAPP-Wild type which is cleaved by β -secretase. Non-crossreacts with sAPP α and full-length APP
- Reference** : 1. Citron M, Oltersdorf T, Haass C, McConlogue L, Hung AY, Seubert P, Vigo-Pelfrey C, Lieberburg I, Selkoe DJ. Mutation of the beta-amyloid precursor protein in familial Alzheimer's disease increases beta-protein production. *Nature* **360** (6405): 672-674 (1992)
2. Goate A, Chartier-Harlin MC, Mullan M, et al. Segregation of a missense mutation in the amyloid precursor protein gene with familial Alzheimer's disease. *Nature* **349** 704-706 (1991)
3. Selkoe DJ. Normal and abnormal biology of the β -amyloid precursor protein. *Annu. Rev. Neurosci.* **17** 489-517 (1994)

For Non-Clinical Research Use Only

IBL Co., LTD. 5-1 Aramachi, Takasaki-Shi, Gunma, 370-0831, JAPAN
TEL: +81 (0)27-310-8040 FAX: +81 (0)27-310-8045 URL: <http://www.ibl-japan.co.jp> E-Mail: do-ibl@ibl-japan.co.jp