

Code No. 18973

**Anti-  
Rab27B Rabbit IgG Affinity Purify**

Volume : 100 uG

- 
- Introduction** : Small GTPase Rab is a large family of membrane trafficking proteins that are conserved in all eukaryotic cells. More than 60 Rab isoforms have been reported in mice and humans, and they are believed to regulate various steps (or various types) of organelle transport.  
The Rab27 subfamily is phylogenetically similar to the Rab3 subfamily, and two Rab27 isoforms, Rab27A and Rab27B, are present in mice and humans. Mutations in the RAB27A gene cause human Griscelli syndrome, which is characterized by pigment dilution and immunodeficiency. The GTP-bound activated form of Rab27A regulates melanosome transport in melanocytes and secretory granule exocytosis (e.g., insulin secretion) through interaction with a specific effector molecule (e.g., Synaptotagmin-like protein (Slp) and Slac2). Rab27B is also expressed on secretory granules, the same as Rab27A, and regulates their exocytosis (e.g., amylase release from rat parotid acinar cells).
- Antigen** : Recombinant Rab27B protein
- Source** : Rabbit antiserum
- Purification** : Affinity Purified with antigen
- Form** : Lyophilized product from 1% BSA in PBS containing 0.05% NaN<sub>3</sub>
- How to use** : 1.0 ml distilled water will be added to the product, then its concentration comes to 100 ug/ml
- Stability** : Lyophilized product, 5 years at 2 – 8 °C  
: Solution, 2 years at –20 °C
- Application** : This antibody can be used for western blotting in concentration of about 1~5  $\mu$ g/mL.
- Specificity** : Cross-reacts with Mouse and Rat.  
Non cross-reacts to Rab3 subfamily
- Reference** : 1. Ménasché, G., Pastural, E., Feldmann, J., Certain, S., Ersoy, F., Dupuis, S., Wulffraat, N., Bianchi, D., Fischer, A., Le Deist, F., de Saint Basile, G. Mutations in *RAB27A* cause Griscelli syndrome associated with haemophagocytic syndrome. *Nat. Genet.* 25:173-176 (2000)  
2. Imai, A., Yoshie, S., Nashida, T., Shimomura, H., Fukuda, M. The small GTPase Rab27B regulates amylase release from rat parotid acinar cells. *J. Cell Sci.* 117:1945-1953 (2004)  
3. Fukuda, M. Versatile role of Rab27 in membrane trafficking: Focus on the Rab27 effector families. *J. Biochem.* 137:9-16 (2005)
- 

For research use only, not for use in diagnostic procedures.

Distributed by:



Immuno-Biological Laboratories, Inc. Toll-Free: 888-523-1246  
8201 Central Ave NE, Suite P Email: info@IBL-America.com  
Minneapolis, MN 55432 Web: www.IBL-America.com