

Code No. 28041

Anti-

GZF1 Rabbit IgG Affinity Purify

Volume : 100 µg

Introduction: GZF1 has been identified as a gene whose expression is induced by a neurotrophic

factor, GDNF. It codes a protein which has BTB/POZ domain in its N-terminal region and 10 of zinc finger motifs in its C-terminal area from the center. It has been known by performance analysis that GZF1 is a transcriptional repression factor sequence-specifically binds to DNA and the binding sequence is present in

transcription control region of HOXA10.

Antigen : Synthetic peptide of the part of Human GZF1

Purification: Purified with antigen peptide

: Lyophilized product from 1% BSA in PBS containing 0.05% NaN₃ Form

How to use : 1.0 mL deionized water will be added to the product (the conc. comes up 100 μg /mL)

: Lyophilized product, 5 years at 2 - 8 °C Stability

: Solution, 2 years at -20 °C

Application: This antibody can be used for western blotting in concentration of about 0.1 µg/mL.

: This antibody can be used for immuno-precipitation in concentration about 1 µg /mL.

Specificity : React with human and mouse GZF1

: 1. Fukuda N, Ichihara M, Morinaga T, Kawai K, Hayashi H, Murakumo Y, Matsuo Reference

> S, Takahashi M. Identification of a novel glial cell line-derived neurotrophic factor-inducible gene required for renal branching morphogenesis. J Biol Chem.

2003 Dec 12;278(50):50386-92.

2. Morinaga T, Enomoto A, Shimono Y, Hirose F, Fukuda N, Dambara A, Jijiwa M, Kawai K, Hashimoto K, Ichihara M, Asai N, Murakumo Y, Matsuo S, Takahashi

M. GDNF-inducible zinc finger protein 1 is a sequence-specific transcriptional repressor that binds to the HOXA10 gene regulatory region. Nucleic Acids Res.

2005 Jul 26;33(13):4191-201.



Email: info@IBL-America.com Web: www.IBL-America.com