

Code No. 28153

**Anti-Human  
LRAT (Lecithin Retinol Acyltransferase) (031) Rabbit IgG A.P. (Affinity Purify)**Volume : 50 µg

---

**Introduction** : LRAT is a retinol-esterifying enzyme and it is mainly found in G0 stage of stellate cells (Ito cells) in liver. At stellate cells, LRAT is thought to take a role engulfing and pooling excess vitamin A released from hepatocytes. As normal hepatocytes are characterized by lipid droplets pooling esterified vitamin A, LRAT is expected to be a marker of stellate cells in G0 stage. LRAT is also expressed in retinal pigment epithelium, small intestine epithelium and skin keratinocyte as well as liver. The physiological significance of LRAT protein at the endothelium is not clarified, yet. This antibody reacts with stellate cells and endothelia of human.

**Antigen** : Synthetic peptide of a part of Human LRAT (GKDKGRNSFYETSSF)

**Purification** : Purified with antigen peptide

**Form** : Lyophilized product from 1 % BSA in PBS containing 0.05 % NaN<sub>3</sub>

**How to use** : 1.0 mL deionized water will be added to the product (the conc. comes up 50 µg /mL)

**Stability** : Lyophilized product, 5 years at 2 – 8 °C  
: Solution, 2 years at –20 °C

**Application** : This antibody can be used for immunohistochemistry with formalin fixed paraffin embedded tissues after microwave treatment or untreated, by several techniques such as Avidin Biotin Complex (ABC) Method. The optimal concentration is about 1 - 5 µg/mL, however, the concentration should be optimized by each laboratory.  
: This antibody can be used for western blotting in concentration of 1 - 5 µg /mL.

**Specificity** : React with Human LRAT.

**Reference** : 1. Nagatsuma K, Hano H, Murakami K, Shindo D, Matsumoto T, Mitobe J, Tanaka K, Saito M, Maehashi H, Owada M, Ikegami M, Tsubota A, Ohkusa T, Aizawa Y, Takagi I, Tajiri H, Matsuura T. Hepatic stellate cells that coexpress LRAT and CRBP-1 partially contribute to portal fibrogenesis in patients with human viral hepatitis. *Liver Int.* 2014 Feb; 34(2):243-52.  
2. Nagatsuma K, Hayashi Y, Hano H, Sagara H, Murakami K, Saito M, Masaki T, Lu T, Tanaka M, Enzan H, Aizawa Y, Tajiri H, Matsuura T. Lecithin: retinol acyltransferase protein is distributed in both hepatic stellate cells and endothelial cells of normal rodent and human liver. *Liver Int.* 2009 Jan; 29(1):47-54.

---

*For research use only, not for use in diagnostic procedures.*

Distributed by:



Immuno-Biological Laboratories, Inc. Toll-Free: 888-523-1246  
8201 Central Ave NE, Suite P Email: info@IBL-America.com  
Minneapolis, MN 55432 Web: www.IBL-America.com