Product Information

Recombinant human Aflatoxin B1 aldehyde

reductase member 3/AKR7A3 protein

Catalog Number: IBATGP0476

PRODUCT INPORMATION

Expression system E.coli

Domain 1-331aa

UniProt No. 095154

NCBI Accession No. AAH25709.1

Alternative Names

Aldo-keto reductase family 7 member A 3, Aflatoxin B1 aldehyde reductase member 3, AFB1 aldehyde reductase 2(AFAR2), Aldo-keto reductase family 7, member A 3 AFAR 2, AFAR2, AFB1 AR 2, Aflatoxin aldehyde reductase, Aflatoxin B1 aldehyde reductase 2, Aflatoxin B1 aldehyde reductase member 3, Aldo keto reductase family 7 member A3, OTTHuMP0000002623.

PRODUCT SPECIFICATION

Molecular Weight

41.6 kDa (370aa) confirmed by MALDI-TOF

Concentration

0.5mg/ml (determined by Bradford assay)

Formulation

Liquid. In Phosphate-Buffered Saline (pH 7.4) containing 10% glycerol

Purity

> 95% by SDS-PAGE

Biological Activity

Specific activity is > 700pmol/min/ug, and is defined as the amount of enzyme that catalyze the reduction 1.0pmole of 1,2-Naphthoquinone presence of NADPH per minute at pH 7.0 at 25C.

Tag

His-Tag

Application SDS-PAGE, Enzyme Activity

Storage Condition

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

For research use only. This product is not intended or approved for human, diagnostics or veterinary use.







Product Information

Recombinant human Aflatoxin B1 aldehyde

reductase member 3/AKR7A3 protein

Catalog Number: IBATGP0476

Description

Aldo-keto reductase family 7, member A 3, also known as AKR7A3, is a member of the aldo/keto reductase superfamily, which consists of more than 40 known enzymes and proteins. This member includes a number of related monomeric NADPH-dependent oxidoreductases, such as aldehyde reductase, aldose reductase, prostaglandin F synthase, xylose reductase, rho crystallin, and many others. AKR7A3 is involved in the detoxification of aldehydes and ketones. The activity of AKR7A3 may detoxify the aflatoxin B1 (AFB1) dialdehyde, which reacts with proteins, and thereby inhibits AFB 1 induced toxicity. Recombinant human AKR7A3, fused to His-tag at N-terminus, was expressed in E. coli and purified by using conventional chromatography techniques.

Amino acid Sequence

<MRGSHHHHHH GMASMTGGQQ MGRDLYDDDD KDRWGSELE>M SRQLSRARPA TVLGAMEMGR RMDAPTSAAV TRAFLERGHT EIDTAFVYSE GQSETILGGL GLRLGGSDCR VKIDTKAIPL FGNSLKPDSL RFQLETSLKR LQCPRVDLFY LHMPDHSTPV EETLRACHQL HQEGKFVELG LSNYAAWEVA EICTLCKSNG WILPTVYQGM YNAITRQVET ELFPCLRHFG LRFYAFNPLA GGLLTGKYKY EDKDGKQPVG RFFGNTWAEM YRNRYWKEHH FEGIALVEKA LQAAYGASAP SMTSATLRWM YHHSQLQGAH GDAVILGMSS LEQLEQNLAA AEEGPLEPAV VDAFNQAWHL VAHECPNYFR

General References

Kniqht LP., et al. (1999) Carcinogenesis. 20(7):1215-23. Bodreddiqari S., et al. (2008) Chem Res Toxicol. 21(5):1134-42.

DATA

SDS-PAGE



3ug by SDS-PAGE under reducing condition and visualized by coomassie blue stain.

For research use only. This product is not intended or approved for human, diagnostics or veterinary use.



